

THE EFFECT OF SIDE STEP ON LOWER EXTREMITY BIOMECHANICS DURING BLOCK LANDING IN FEMALE VOLLEYBALL PLAYERS

Chen-Fu Huang and Yao-Yi Hsieh

Department of Physical Education, National Taiwan Normal University, Taipei, Taiwan

The purpose of this study was to examine the effect of block and side step block on lower extremity biomechanics during landing in female volleyball players. Eight female university volleyball players participated in this study. The kinematic and kinetic data were collected by eight Vicon cameras (250 Hz) and two force plates (1000 Hz). The Visual 3D software was used to analyze the kinematic and kinetic of block and side step block landing variables. The results showed a significantly higher knee extensor moment during side step landing than the block landing at the time of peak vertical ground reaction force and peak joint moment. It was concluded that female players displayed greater knee extensor moment during the side step before block landing that may increase the loading on the knee.

KEY WORDS: injury, lateral movement, joint moment, sagittal plane.

INTRODUCTION: Block is the first line of defense and an offensive skill in the volleyball competition. Many volleyball players perform the lateral movement before the block. The volleyball competition showed that players jump over 80% of the time after lateral movement (Lobiatti, 2008). Although, volleyball competition is a non-contact sport, but it had a high musculoskeletal injury rate after landing movement (Briner & Kacmar, 1997). Many volleyball players often occurred injuries in lower extremity after landing movement (Gerberich, 1987). ACL re-rupture is a serious lower extremity injury, and landing is one of the primary mechanisms for ACL injury (Ferretti, Papandrea, Conteduca, & Mariani, 1992). Previous studies (Arendt & Dick, 1995; Ferretti et al., 1992) reported that the females had non-contact ACL injury rate four to eight times greater than males in the same competitions such as basketball, handball, and volleyball. Several studies indicated that females performed landing movement with more erect posture (Colby et al., 2000; Lephart et al., 2002; Rozzi, Lephart, Gear, & Fu, 1999; Schmitz, Kulas, Perrin, & Riemann, 2007), and exhibited greater knee extensor moment (Salci et al., 2004; Hughes, Watkins, & Owen, 2010) that produced a great loading on the knee joint. Therefore, it was important to reduce the ground impact more efficiently during landing. Hughes et al. (2010) reported that female displayed greater knee flexion angle than male during volleyball block landing, but it were no differences in the joint moment in the sagittal plane. Previous study (Hughes et al., 2008, 2010) mainly focused on block landing, but players mostly performed lateral movement before block landing. To our knowledge, no study investigates the effect of lateral movement on lower extremity biomechanics during volleyball block landing. The purpose of this study was to examine the effect of stand and side step on lower extremity biomechanics during block and side step block landing in female volleyball players, and to understand the important variable of block and side step block landing movement in the injury risk of lower extremity.

METHODS: Eight females (age: 20.4±1 yrs, height: 170.3±5 cm, weight: 63.6±6.2 kg) were recruited from the university volleyball team. All participants signed informed consent before the study. Two adjacent force plates (Kistler 9287 & AMTI 5507) embedded into the floor sampling at 1000 Hz were used to measure ground reaction force to determine initial ground contact of right and left legs on landing. An 8-camera Vicon system (Vicon T20-S, Oxford, UK), sampling at 250 Hz, was used to collect the three-dimensional (3D) coordinates of 51 retro-reflective markers. Markers were placed directly on the skin of 15 segments for each participants. These segments include: head, trunk, right and left upper arm, right and left

Ankle dorsiflexion (+)	-28.9 (3.5)	-29.9 (3.2)
PVGRF joint angle (°)		
Hip flexion (+)	13.9 (6.9)	10.9 (7.7)
Knee flexion (+)	45.5 (5.3)	46.4 (5.9)
Ankle dorsiflexion (+)	26.6 (1.6)	27.8 (2.4)
Maximum joint angle(°)		
Hip flexion (+)	24.6 (14.3)	23.1 (17.6)
Knee flexion (+)	62.5 (11.6)	64.3 (11.9)
Ankle dorsiflexion (+)	31.5 (2.6)	32.2 (2.7)
Range of motion (°)		
Hip	23.3 (11.7)	24.2 (14.9)
Knee	51.6 (12.3)	52 (11.9)
Ankle	60.4 (1.9)	62.2 (1.7)

*Significant difference between block and side step block ($P<0.05$); IC = Initial contact; PVGRF = at the time of peak vertical ground reaction force. Positive values indicate flexion angle and negative values indicate extension angle.

Table 2

Means(SD) of the vertical ground reaction force and joint moments between block and side step block landing

Kinetic variables	Block jump	Side step block
VGRF (BW)	2.89 (0.82)	3.33 (0.77)
IC joint moment (Nm/kg)		
Hip extensor	-0.42 (0.36)	-0.47 (0.27)
Knee extensor	-0.21 (0.20)	-0.24 (0.17)
Ankle plantar flexor	-0.05 (0.07)	-0.05 (0.10)
PVGRF joint moment (Nm/kg)		
Hip extensor	-0.87 (0.78)	-1.06 (0.76)
Knee extensor *	-2.51 (0.86)	-3.08 (0.82)
Ankle plantar flexor	-2.28 (0.71)	-2.27 (0.52)
Peak joint moment (Nm/kg)		
Hip extensor	-2.04 (0.52)	-1.86 (0.44)
Knee extensor *	-3.03 (0.91)	-3.59 (0.81)
Ankle plantar flexor	-2.52 (0.77)	-2.52 (0.54)

*Significant difference between block and side step block ($P<0.05$); IC = Initial contact; PVGRF = at the time of peak vertical ground reaction force. Positive values indicate flexor moment and negative values indicate extensor moment.

DISCUSSION: The result showed that there were no significantly lower extremity joint angles between block jump landing and side step block landing. Although, in this study there is no difference in the kinematics between block jump landing and side step block landing, but we found that there were differences in the knee joint moment during landing when female performed side step block landing. The result showed that a greater knee extensor moment during side step landing at the time of peak vertical ground reaction force. It was indicated that knee joint had a greater loading during side step block landing. Chappell et al. (2002) reported that the landing phase generates larger anterior knee shear forces compared with the takeoff phase. And females exhibit greater proximal tibia anterior shear forces and larger knee extension during the landing phase. We also found that a greater peak knee extensor moment during side step landing. A number of studies (Salci et al., 2004; Yu et al., 2006; Hughes et al., 2010) reported that females exhibited a greater knee extensor moment during

landing, it may increase the risk of ACL injury. A greater eccentric quadriceps contraction that was increased knee extension moment results in higher external ground reaction forces, which increased anterior translation of the tibia relative to the femur, that might occur ACL injury (Ball et al., 1999). Therefore, females performed lateral side step movement before the block increase loading on knee joint.

CONCLUSION: This study demonstrated that lateral movement may alter lower extremity biomechanics during landing. It was concluded that female players displayed greater knee extensor moment during the side step before block landing that may increase the loading on the knee. It was suggested that volleyball coaches and players to decrease lateral side step movement before block landing while lower extremity muscles fatigue in the training. That might reduce overloading on the knee joint and risk of injury.

REFERENCES:

- Arendt, E., & Dick, R., (1995). Knee injury patterns among men and women in collegiate basketball and soccer. NCAA data and review of literature. *American Journal of Sports Medicine*, 23(6), 694–701.
- Ball, K. A., Brooks-Hill, A. L., Richards, D., Mark, P., & Evans, R. J. (1999). Lack of hip flexion: A mechanism for ACL injury. *Medicine & Science in Sport & Exercise*. 31(5), S295.
- Briner, W. W., & Kacmar, L. (1997). Common injuries in volleyball. *Sports Medicine*, 24(1), 67–71.
- Chappell, J. D., Yu, B., Kirkendall, D. T., & Garrett, W. E. (2002). A comparison of knee kinetics between male and female recreational athletes in stop-jump tasks. *The American Journal of Sports Medicine*. 30, 261-267.
- Colby, S., Francisco, A., Yu, B., Kirkendall, D., Finch, M., & Garrett, W. (2000). Electromyographic and kinematic analysis of cutting maneuvers. Implications for anterior cruciate ligament injury. *American Journal of Sports Medicine*. 28(2), 234-240.
- Ferretti, A., Papandrea, P., Conteduca, F., & Mariani P. P. (1992). Knee ligament injuries in volleyball players. *The American Journal of Sports Medicine*, 20(2), 203-207.
- Gerberich, S. G. (1987). Analysis of severe injury associated with volleyball activities. *Physical and Sports Medicine*, 15(8), 75-79.
- Hughes, G., Watkins, J., & Owen, N. (2008). Gender differences in lower limb frontal plane kinematics. *Sport Biomechanics*, 7(3), 333-341.
- Hughes, G., Watkins, J., & Owen, N. (2010). Differences between the sexes in knee kinetics during landing from volleyball block jumps. *European Journal of Sport Science*, 10(1), 1-11.
- Lephart, S. M., Ferris, C. M., Riemann, B. L., Myers, J. B., & Fu, F. H. (2002). Gender Differences in Strength and Lower Extremity Kinematics During Landing. *Clinical Orthopaedics & Related Research*, 140, 162-169.
- Lobietti, R., Fantozzi, S., Stagni, R., & Merni, F. (2008). Kinematics analysis of landing from volleyball block. *Gait & Posture*, 28(1), 19.
- Rozzi, S.L., Lephart, S. M., Gear, W. S., & Fu, F. H. (1999). Knee joint laxity and neuromuscular characteristics of male and female soccer and basketball players. *The American Journal of Sports Medicine*. 27(3), 312–319.
- Salci, Y., Kentel, B. B., Heycan, C., Akin, S., & Korkusuz, F. (2004). Comparison of landing maneuvers between male and female college volleyball players. *Clinical Biomechanics*, 19, 622-628.
- Schmitz, R. J., Kulas, A. S., Perrin, D. H., & Riemann, B. L. (2007). Sex differences in lower extremity biomechanics during single leg landings. *Clinical Biomechanics*, 22(6), 681-688.
- Yu, B., Lin, C. F., & Garrett, W. E. (2006). Lower extremity biomechanics during the landing of a stop-jump task. *Clinical Biomechanics*. 21, 297-305.

Acknowledgement

This study was funded by a grant from National Science Council (NSC 101-2410-H-003-129) in Taiwan.